

41st INTERNATIONAL CHEMISTRY OLYMPIAD

UK Round One - 2009

MARKING SCHEME

Notes

Chemical equations may be given as sensible multiples of those given here.
Formulae can be given by any conventional method (i.e. structural or molecular).

State symbols do not need to be included in the chemical equations to obtain the mark(s).

Answers should be given to an appropriate number of significant figures
although the marker should only penalise this once.

Total 64 marks.

Question 1		Answer	Marks
(a)	i)	It is aromatic / the bonds in the ring are conjugated / there are alternate single and double bonds in the ring / the electrons in the ring are delocalised / very similar to benzene	1
	ii)	3 peaks	1
	iii)	$C_2H_6N_8 + 7/2 O_2 \rightarrow 2 CO_2 + 3 H_2O + 4 N_2$ or $2 C_2H_6N_8 + 7 O_2 \rightarrow 4 CO_2 + 6 H_2O + 8 N_2$	1
(b)	i)	19° or 19.5° or 19°28'	1
	ii)	$C_8N_8O_{16} \rightarrow 8 CO_2 + 4 N_2$	1
(c)		$C_6H_7N_3O_{11} + 9/4 O_2 \rightarrow 6 CO_2 + 7/2 H_2O + 3/2 N_2$ or $4 C_6H_7N_3O_{11} + 9 O_2 \rightarrow 24 CO_2 + 14 H_2O + 6 N_2$	2

7 marks


Note: Tests are to be taken under controlled conditions. Students must not have access to the information contained in this marking scheme prior to, or during, the test.

Question 2			
		Answer	Marks
(a)	i)	Moles = $13000/44.1 = 295$	1
	ii)	Mass = $3 \times 295 \times 44.0 = 38900 \text{ g} = 38.9 \text{ kg}$ (accept 39 kg)	1
	iii)	Heat energy = $2220 \times 295 = 655000 \text{ kJ} = 655 \text{ MJ}$	1
	iv)	$1 \text{ mol s}^{-1} = 2220 \text{ kJ s}^{-1} = 2220 \text{ kW}$, so 15 kW $= 15/2220 \text{ mole s}^{-1} = 15 \times 24000/2220 = 162 \text{ cm}^3 \text{ s}^{-1}$	1
	v)	Still 140 psi (or 9.52 atm)	1
(b)	i)	Sensible bonding diagram with all single covalent bonds Accept a bond angle anything between $90^\circ - 105.5^\circ$	1 1
	ii)	Mass = $295 \times 0.02 \times 10^{-9} \times 62.1 = 0.000000366 \text{ g}$ $= 0.000366 \text{ mg}$ $= 3.66 \times 10^{-7} \text{ g}$ (accept 3.7 or $4.0 \times 10^{-7} \text{ g}$)	1
(c)		$6000 \text{ m}^3 \text{ CH}_4 = 6000 \times 10^3 \text{ dm}^3 = 6000 \times 10^3 / 24 \text{ moles}$, so we get $6000 \times 10^3 / 24 \text{ moles CO}_2 = (6000 \times 10^3 / 24) \times 44 \text{ g CO}_2$ per hour. So in 16 days we get $(6000 \times 10^3 / 24) \times 44 \times 24 \times 16 = 4224 \times 10^6 \text{ g} = 4224 \text{ tonnes}$ (accept 4200 tonnes)	1





9 marks

Question 3			
		Answer	Marks
(a)	i)	$\text{P}_4\text{S}_3 + 8\text{O}_2 \rightarrow \text{P}_4\text{O}_{10} + 3\text{SO}_2$ (accept $2\text{P}_2\text{O}_5 + 3\text{SO}_2$)	1
	ii)	$2\text{KClO}_3 \rightarrow 2\text{KCl} + 3\text{O}_2$	1
	iii)	$3\text{P}_4\text{S}_3 + 16\text{KClO}_3 \rightarrow 3\text{P}_4\text{O}_{10} + 9\text{SO}_2 + 16\text{KCl}$ (accept $6\text{P}_2\text{O}_5$)	1
	iv)	$\text{P}_4\text{S}_3 / \text{KClO}_3 = 660 / 1961 = 1 / 2.97$	1





Note: Tests are to be taken under controlled conditions. Students must not have access to the information contained in this marking scheme prior to, or during, the test.

	v)	$\Delta_r H^\ominus = (3 \times -2948) + (9 \times -296.8) + (16 \times -436.7) - (3 \times -154.0) + (16 \times -397.7)$ $= -11700 \text{ kJ mol}^{-1}$	2
(b)	i)	3 peaks	1
	ii)	4 peaks	1
	iii)	3 peaks	1
(c)		 <p>(accept any other reasonable structure, that fits with the data and with elements in correct valencies)</p>	2


11 marks

Question 4		Answer	Marks
(a)		 <p>Loperamide</p>	1
(b)	i)	 <p>carboxylic acid</p>  <p>alcohol</p>	1
	ii)		1

Note: Tests are to be taken under controlled conditions. Students must not have access to the information contained in this marking scheme prior to, or during, the test.

(c)		 <p>anion C cyclic ester D</p>	2
(d)		 <p>E F</p>	2
(e)	i)	 <p>open-chain amide</p>	1
	ii)	 <p>cyclic bromide salt</p>	2

10 marks

Question 5		Answer	Marks
(a)	i)	It must be clear from the structure that arsine is not planar. Structures similar to those shown below would be acceptable: 	1
	ii)	$4\text{AsH}_3 + 3\text{O}_2 \rightarrow 4\text{As} + 6\text{H}_2\text{O}$	1
(b)	i)	As_2O_3	1
	ii)	$\text{As}_2\text{O}_3 + 6\text{Zn} + 6\text{H}_2\text{SO}_4 \rightarrow 2\text{AsH}_3 + 6\text{ZnSO}_4 + 3\text{H}_2\text{O}$	1
(c)	i)	+5 or (V)	1

Note: Tests are to be taken under controlled conditions. Students must not have access to the information contained in this marking scheme prior to, or during, the test.

	ii)	Both the bonding and geometry must be clear, structures such as those shown below would be acceptable:	1
(d)	i)	From the graph the $t_{1/2}$ is 8 mins $k = \ln 2 / t_{1/2}$ therefore $k = 0.087 \text{ min}^{-1}$ (0.0014 s^{-1} or $0.144 \times 10^{-3} \text{ s}^{-1}$) Accept values for $t_{1/2}$ in the region of 7 to 9 mins ($k = 0.08$ to 0.1 min^{-1}) also accept correct values for k given in s^{-1} .	1
	ii)	$[\text{HAsO}_4^{2-}(\text{aq})]_t = [\text{HAsO}_4^{2-}(\text{aq})]_0 \exp^{-kt}$ $10 = [\text{HAsO}_4^{2-}(\text{aq})]_0 \exp^{-(0.09 \times 55)}$ $[\text{HAsO}_4^{2-}(\text{aq})]_0 = 1400 \mu\text{g dm}^{-3}$ For $k = 0.08 \text{ min}^{-1}$, $[\text{HAsO}_4^{2-}(\text{aq})]_0 = 800 \mu\text{g dm}^{-3}$ whilst for $k = 0.1 \text{ min}^{-1}$, $[\text{HAsO}_4^{2-}(\text{aq})]_0 = 2400 \mu\text{g dm}^{-3}$. Full marks should be given for values within this range.	2
(e)		$[\text{HAsO}_4^{2-}(\text{aq})]_{t=0} = [\text{HAsO}_4^{2-}(\text{aq})]_{\text{eq}} + [\text{HAsO}_4^{2-}(\text{adsorbed})]_{\text{eq}}$ Therefore: $K = \frac{[\text{HAsO}_4^{2-}(\text{aq})]_{t=0} - [\text{HAsO}_4^{2-}(\text{aq})]_{\text{eq}}}{[\text{HAsO}_4^{2-}(\text{aq})]_{\text{eq}}}$ Rearranges to give: $[\text{HAsO}_4^{2-}(\text{aq})]_{\text{eq}} = \frac{[\text{HAsO}_4^{2-}(\text{aq})]_{t=0}}{1 + K} = \frac{30}{1 + 186} = 0.16 \mu\text{g/dm}^3$	2

11 marks

Question 6			
		Answer	Marks
(a)	i)	-13.6 eV Must have minus sign	1
	ii)	zero	1
	iii)	1300 kJ mol ⁻¹	1
(b)		$-495.8 \times 10^3 = -1312 \times 10^3 \times \frac{Z_{\text{eff}}^2}{3^2}$ $Z_{\text{eff}} = 1.84$ (2 for correct answer; partial credit of 1 if expression is correct)	2

Note: Tests are to be taken under controlled conditions. Students must not have access to the information contained in this marking scheme prior to, or during, the test.

(c)		<p>energy of electron in 2p shell = $-13.6 \times \frac{(Z - S)^2}{2^2}$</p> <p>energy of electron in 1s shell = $-13.6 \times \frac{(Z - S)^2}{1^2}$</p> <p>energy released on transition from 2p to 1s</p> <p>$= \frac{3}{4} \times 13.6 \times (Z - 1)^2 = 8000$</p> <p>$(Z - 1) = 28$</p> <p>$Z = 29$ element is copper, Cu</p>	2
(d)	i)	<p>energy of electron in 3d shell = $-13.6 \times \frac{(Z - S)^2}{3^2}$</p> <p>energy of electron in 2p shell = $-13.6 \times \frac{(Z - S)^2}{2^2}$</p> <p>energy released on transition from 3d to 2p</p> <p>$= \left(\frac{1}{2^2} - \frac{1}{3^2} \right) \times 13.6 \times (Z - 7.4)^2 = 10000$</p> <p>$(Z - 7.4) = 72.8$</p> <p>$Z = 80$ element is mercury, Hg</p>	2
	ii)	HgS (accept other possible mercury sulfide formulae)	1
(e)	i)	<p>for C, energy released on transition from 2p to 1s</p> <p>$= \frac{3}{4} \times 13.6 \times (Z - 1)^2 = 10500$</p> <p>$(Z - 1) = 32$</p> <p>$Z = 33$ element is arsenic, As</p> <p>for D energy released on transition from 3d to 2p</p> <p>$= \left(\frac{1}{2^2} - \frac{1}{3^2} \right) \times 13.6 \times (Z - 7.4)^2 = 10500$</p> <p>$(Z - 7.4) = 74.6$</p> <p>$Z = 82$ element is lead, Pb</p>	1 1
	ii)	CuAsHO ₃ (this assumes +2 oxidation state for Cu)	2
(f)		<p>for antimony, Sb, energy released on transition from 2p to 1s</p> <p>$= \frac{3}{4} \times 13.6 \times (51 - 1)^2 = 25500 \text{ eV}$</p>	1
(g)		<p>Balancing oxidation states:</p> <p>$(2x) + (2 \times 5) + (7 \times -2) = 0$ implies $x = +2$</p> <p>[formula is Pb₂Sb₂O₇]</p>	1

16 marks

Total Marks 64

Note: Tests are to be taken under controlled conditions. Students must not have access to the information contained in this marking scheme prior to, or during, the test.